

Partial recovery of entanglement in bipartite entanglement transformations*

Somshubhro Bandyopadhyay¹, Vwani Roychowdhury¹, Farrokh Vatan^{1,2}

1. Electrical Engineering Department, UCLA, Los Angeles, CA 90095

2. Jet Propulsion Laboratory, California Institute of Technology 4800 Oak Grove Drive Pasadena, CA 91109

Abstract

Any *deterministic* bipartite entanglement transformation involving finite copies of pure states and carried out using local operations and classical communication (LOCC) results in a net loss of entanglement. We show that for almost all such transformations, partial recovery of lost entanglement is achievable by using 2×2 auxiliary entangled states, no matter how large the dimensions of the parent states are. For the rest of the special cases of deterministic LOCC transformations, we show that the dimension of the auxiliary entangled state depends on the presence of equalities in the majorization relations of the parent states. We show that genuine recovery is still possible using auxiliary states in dimensions less than that of the parent states for *all* patterns of majorization relations except only one special case.

Entanglement, shared among spatially separated parties, is a critical resource that enables efficient implementations of several quantum information processing [2] and distributed computation [3] tasks. To better exploit the power of entanglement, considerable effort has been put into understanding its transformation properties [4]–[7] and characterizing transformations allowed under local operations and classical communication (LOCC). A central question is: what happens to the overall entanglement during transformations? In the asymptotic limit involving infinite number of copies of pure states, entanglement can be

*This work was sponsored in part by the Defense Advanced Research Projects Agency (DARPA) project MDA972–99–1–0017 [note that the content of this paper does not necessarily reflect the position or the policy of the government, and no official endorsement should be inferred], and in part by the U.S. Army Research Office/DARPA under contract/grant number DAAD19–00–1–0172.

The research of Farrokh Vatan was performed partly at UCLA and supported by the above mentioned grants and was performed partly at the Jet Propulsion Laboratory (JPL), California Institute of Technology, under contract with National Aeronautics and Space Administration (NASA). The Revolutionary computing Technologies Program of the JPL’s Center for Integrated Space Microsystems (CISM) supported his work.

Emails: som@ee.ucla.edu, vwani@ee.ucla.edu, Farrokh.Vatan@jpl.nasa.gov

concentrated and diluted with unit efficiency [4]. This remarkable asymptotic "nondissipative" property, however, does not hold in the finite copy regime, where the process becomes inherently "dissipative," and a local *deterministic* conversion between two pure entangled states (which are not locally unitarily related), *always* results in a net loss of entanglement [5].

It is of fundamental importance to devise local strategies to *recover* the lost entanglement in an entanglement manipulation. Such recovery strategies would require *collective* manipulations with ancillary resources. That is, let $|\psi\rangle = \sum_{i=1}^n \sqrt{\alpha_i} |i\rangle |i\rangle$ and $|\varphi\rangle = \sum_{i=1}^n \sqrt{\beta_i} |i\rangle |i\rangle$ be, respectively, the source and target states in $n \times n$ such that $|\psi\rangle \rightarrow |\varphi\rangle$ under LOCC. Then the amount of entanglement lost in such a transformation is $E(|\psi\rangle) - E(|\varphi\rangle)$ (where E is the entropy of entanglement, e.g., $E(|\psi\rangle) = -\sum_{i=1}^n \alpha_i \ln(\alpha_i)$), and we say that there is a *partial recovery* of the lost entanglement if there exist entangled states $|\chi\rangle, |\omega\rangle$ in $k \times k$, $k < n$, such that $|\psi\rangle \otimes |\chi\rangle \rightarrow |\varphi\rangle \otimes |\omega\rangle$ with certainty under LOCC, and $E(|\omega\rangle) > E(|\chi\rangle)$. Since the overall transformation involving the auxiliary states is dissipative, the recovered entanglement, $(E(|\omega\rangle) - E(|\chi\rangle))$, is always less than or equal to the initial amount of lost entanglement, $(E(|\psi\rangle) - E(|\varphi\rangle))$. In order to minimize the use of ancillary resources and to reduce the complexity of the collective operations, we consider a partial recovery of entanglement process to be *efficient*, if the dimension of the auxiliary states, k , is the minimum required for the recovery process to happen. Moreover, in order for the partial recovery process to be *genuine*, we require the dimension of the auxiliary states to be smaller than that of the parent states (i.e., $k < n$), since otherwise, if $k = n$ then one can have a complete recovery of lost entanglement by a trivial choice: $|\chi\rangle = |\varphi\rangle$ and $|\omega\rangle = |\psi\rangle$.

A first step toward achieving partial recovery of entanglement has recently been taken in [8] for the *special case of* $n = 2$. This result is of limited interest only: since the auxiliary pure states are necessarily of the same dimension as the parent states (i.e., $k = n = 2$), one can always have a complete recovery of lost entanglement by a trivial choice of the auxiliary states. However, [8] presents non-trivial selections of auxiliary states (i.e., $|\chi\rangle \neq |\varphi\rangle$) that lead to partial recovery of entanglement.

We prove that genuine and efficient partial recovery of entanglement is *always* possible for almost all bipartite entanglement transformations in *any* finite dimension (i.e., for any $n > 2$). Moreover, *for almost all comparable pairs*, such *partial recovery* is achievable by using auxiliary states of *minimum possible dimension*, i.e., $k = 2$, no matter *how large* the dimensions of the parent states are. For the rest of the special cases of comparable parent states, we show that the dimension of the auxiliary entangled state depends on the structure of the majorization relations of the parent states, where the presence of equalities in the majorization relations either in isolation or in blocks determine the dimension of the auxiliary entanglement.

Recall that our parent bipartite pure states are represented as $|\psi\rangle = \sum_{i=1}^n \sqrt{\alpha_i} |i\rangle |i\rangle$ and $|\varphi\rangle = \sum_{i=1}^n \sqrt{\beta_i} |i\rangle |i\rangle$, where $\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_n$ and $\beta_1 \geq \beta_2 \geq \dots \geq \beta_n$, are the Schmidt coefficients. Hence, the eigenvalues of the reduced density matrices $\rho_\psi \equiv \text{Tr}_{B(\text{or } A)} |\psi\rangle_{AB} \langle\psi|$ and $\rho_\varphi \equiv$

$\text{Tr}_{B(\text{or } A)} |\varphi\rangle_{AB} \langle\varphi|$ are $\alpha_1, \alpha_2, \dots, \alpha_n$ and $\beta_1, \beta_2, \dots, \beta_n$, respectively. Define the vector of the eigenvalues as $\lambda_\psi \equiv (\alpha_1, \dots, \alpha_n)$ and $\lambda_\varphi \equiv (\beta_1, \dots, \beta_n)$. Since our parent states are comparable, i.e., $|\psi\rangle \longrightarrow |\varphi\rangle$ with probability one under LOCC, it follows from [5] that λ_ψ is *majorized* by λ_φ , (denoted as $\lambda_\psi \prec \lambda_\varphi$); i.e.,

$$\sum_{i=1}^m \alpha_i \leq \sum_{i=1}^m \beta_i, \quad \text{for every } m = 1, \dots, n-1. \quad (1)$$

Note that both sides equal one for $m = n$.

First we consider the case where λ_ψ is *strictly majorized* by λ_φ , i.e., all the inequalities of the majorization conditions (1) are *strict*, and show that recovery with an auxiliary entangled state in 2×2 is always possible. We represent strict majorization as $\lambda_\psi \sqsubset \lambda_\varphi$. Note that *for a randomly picked pair of comparable states, the majorization inequalities are strict with probability one*. This guarantees that the case where all the majorization inequalities are strict covers *almost all* possible comparable pairs. We first illustrate the basic idea involved in the proof with a simple example.

Example. Consider the states $|\psi\rangle$ and $|\varphi\rangle$ with $\lambda_\psi = (0.4, 0.3, 0.2, 0.1)$, and $\lambda_\varphi = (0.5, 0.3, 0.2, 0)$. Then $\lambda_\psi \sqsubset \lambda_\varphi$. Note that since $|\psi\rangle \longrightarrow |\varphi\rangle$, then for all states $|\chi\rangle$, $|\psi\rangle \otimes |\chi\rangle \longrightarrow |\varphi\rangle \otimes |\chi\rangle$. For partial recovery, our first strategy is to obtain a 2×2 state $|\chi(p)\rangle$ with $\lambda_{\chi(p)} = (p, 1-p)$ and $p \in (0.5, 1)$ such that $\lambda_{\psi \otimes \chi(p)} \sqsubset \lambda_{\varphi \otimes \chi(p)}$. This allows a perturbation of p to $p - \varepsilon$ ($\varepsilon > 0$) on the right hand side of the underlying $2n$ inequalities such that the inequalities are still satisfied after the perturbation. The second requirement is that such a perturbation should preserve the ordering of the Schmidt coefficients, i.e., the ordering of the coefficients is the same in both $|\varphi\rangle \otimes |\chi(p - \varepsilon)\rangle$ and $|\varphi\rangle \otimes |\chi(p)\rangle$. These two facts together would imply that there is an $\varepsilon > 0$ such that $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p - \varepsilon)}$. We now let $|\chi\rangle = |\chi(p)\rangle$ and $|\omega\rangle = |\chi(p - \varepsilon)\rangle$; since $\varepsilon > 0$, $E(|\omega\rangle) > E(|\chi\rangle)$. Choosing $p = 0.8$ we have $\lambda_{\psi \otimes \chi(p)} \sqsubset \lambda_{\varphi \otimes \chi(p)}$; both the order of the coefficients of $|\varphi\rangle \otimes |\chi(0.8)\rangle$, as well as the relation $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p - \varepsilon)}$ hold if $0 < \varepsilon < 0.08$. Thus, we can choose $|\chi\rangle = |\chi(0.8)\rangle$, and $|\omega\rangle = |\chi(0.73)\rangle$. Then $|\psi\rangle \otimes |\chi\rangle \longrightarrow |\varphi\rangle \otimes |\omega\rangle$, where $E(|\omega\rangle) > E(|\chi\rangle)$. ■

Theorem 1 *If λ_ψ is strictly majorized by λ_φ then there are 2×2 states $|\chi\rangle$ and $|\omega\rangle$ such that $|\psi\rangle \otimes |\chi\rangle \longrightarrow |\varphi\rangle \otimes |\omega\rangle$ and $E(|\omega\rangle) > E(|\chi\rangle)$.*

Proof: Let $|\chi(p)\rangle$ be a 2×2 state with $\lambda_{\chi(p)} = (p, 1-p)$, and $\frac{1}{2} < p < 1$. Note that for all values of $p \in (\frac{1}{2}, 1)$, $|\psi\rangle \otimes |\chi(p)\rangle \longrightarrow |\varphi\rangle \otimes |\chi(p)\rangle$. The choice of p determines the orderings of the Schmidt coefficients of $|\psi\rangle \otimes |\chi(p)\rangle$ and $|\varphi\rangle \otimes |\chi(p)\rangle$, and hence the inequalities in $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p)}$. Conversely, one can think in terms of the orderings of the Schmidt coefficients. There is only a finite number (in fact, at most $n!$) of possible individual orderings of the coefficients of $|\psi\rangle \otimes |\chi(p)\rangle$ and $|\varphi\rangle \otimes |\chi(p)\rangle$. For each

such ordering one can determine its feasible set: values of $p \in (\frac{1}{2}, 1)$ for which the ordering is valid. Each nonempty feasible set corresponds to the solution of a set of linear inequalities, and hence, is a union of intervals and discrete points in $(\frac{1}{2}, 1)$. Moreover, the union of the feasible sets of all possible orderings of the coefficients is the interval $(\frac{1}{2}, 1)$. Hence, it follows from simple measure-theoretic arguments that there exists at least one ordering, where the corresponding feasible set includes intervals of nonzero lengths of the form (a, b) , where $\frac{1}{2} < a < b < 1$.

Next, let us restrict p to belong to such a nontrivial feasible set, F . We next show that $\lambda_{\psi \otimes \chi(p)} \sqsubset \lambda_{\varphi \otimes \chi(p)}$ for all values of $p \in F$, *except* at most $2n - 1$ discrete values. Hence, the set of points p where the majorization inequalities are strict and the ordering of Schmidt coefficients is preserved is of non-zero measure, i.e., it includes intervals. If in the majorization relationship of $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p)}$ one of the inequalities (among the $2n - 1$ nontrivial inequalities) is an equality, then we must have

$$p \sum_{j=1}^x \alpha_j + (1-p) \sum_{j=1}^y \alpha_j = p \sum_{j=1}^s \beta_j + (1-p) \sum_{j=1}^t \beta_j, \quad (2)$$

where $x + y = s + t$, $x \geq y$, and $s \geq t$. Equivalently,

$$\left(\sum_{j=1}^x \alpha_j - \sum_{j=1}^y \alpha_j - \sum_{j=1}^s \beta_j + \sum_{j=1}^t \beta_j \right) p = \sum_{j=1}^t \beta_j - \sum_{j=1}^y \alpha_j. \quad (3)$$

There are two cases now: **(i)** Equation (3) determines a value of p , and **(ii)** Equation (3) is an equivalence, and hence, does not determine a value for p . We show that case **(ii)** is impossible: (3) does not determine a value for p , if and only if $\sum_{j=1}^x \alpha_j = \sum_{j=1}^s \beta_j$ and $\sum_{j=1}^y \alpha_j = \sum_{j=1}^t \beta_j$. Since $\lambda_\psi \sqsubset \lambda_\varphi$, it follows that $x > s$ and $y > t$. This *contradicts* the condition $x + y = s + t$. Hence, every potential equality in the majorization relationship $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p)}$ corresponds to a fixed value for p . Since, there are at most $2n - 1$ such nontrivial equalities, there are at most $2n - 1$ values for p for which $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p)}$ is not strict.

Hence, there exist a $p \in F \subseteq (\frac{1}{2}, 1)$ and an $0 < \varepsilon < \frac{1}{2}$ such that $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p-\varepsilon)}$. The proof is completed by setting $|\chi\rangle = |\chi(p)\rangle$ and $|\omega\rangle = |\chi(p-\varepsilon)\rangle$. ■

What happens if λ_ψ is **not** strictly majorized by λ_φ ? We first define $\Delta_{\psi, \varphi}$ as the set of all indices m such that the relation (1) is an equality:

$$\Delta_{\psi, \varphi} = \left\{ m : 1 \leq m \leq n-1, \sum_{i=1}^m \alpha_i = \sum_{i=1}^m \beta_i \right\}.$$

Note that $1 \in \Delta_{\psi, \varphi}$ is equivalent to the case $\alpha_1 = \beta_1$ and $n-1 \in \Delta_{\psi, \varphi}$ is equivalent to the case $\alpha_n = \beta_n$.

We first show that *even in the presence of many patterns of equalities* in the majorization relationship of the parent states, *recovery is still possible* using only 2×2 auxiliary states.

Theorem 2 Suppose that $1 \notin \Delta_{\psi,\varphi}$ (i.e., $\alpha_1 \neq \beta_1$), $n-1 \notin \Delta_{\psi,\varphi}$ (i.e., $\alpha_n \neq \beta_n$), and if $j \in \Delta_{\psi,\varphi}$ then $j+1 \notin \Delta_{\psi,\varphi}$ (i.e., there are no consecutive equalities in the majorization). Then there are 2×2 states $|\chi\rangle$ and $|\omega\rangle$ such that $|\psi\rangle \otimes |\chi\rangle \longrightarrow |\varphi\rangle \otimes |\omega\rangle$ and $E(|\omega\rangle) > E(|\chi\rangle)$.

Proof: We first show that there exists a nonempty interval $I = (\frac{1}{2}, a)$ ($1 > a > \frac{1}{2}$), such that each inequality in the majorization relationship $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p)}$ is either **(i)** a “benign” identity for all $p \in I$; that is, the equality holds even if on the right hand side p is perturbed to $p - \varepsilon$, for any $\varepsilon > 0$, or **(ii)** is a strict inequality, for all $p \in I$, except for at most $2n - 1$ discrete values. Such a majorization, where each inequality is either strict or a benign identity, is represented as $\lambda_{\psi \otimes \chi(p)} \sqsubseteq \lambda_{\varphi \otimes \chi(p)}$. One can then use the simple measure-theoretic arguments introduced in the proof of Theorem 1, and show that there exists an ordering of the Schmidt coefficients of $|\varphi\rangle \otimes |\chi(p)\rangle$ such that $F \cap I$ has a nonzero measure (i.e., includes intervals), where F is the feasible set for the given ordering. The two above results then show that there exists a $p \in F \cap I$ such that $\lambda_{\psi \otimes \chi(p)} \sqsubseteq \lambda_{\varphi \otimes \chi(p)}$, and in a neighborhood around p the ordering of the Schmidt coefficients of $|\varphi\rangle \otimes |\chi(p)\rangle$ is preserved. Hence, there is an $0 < \varepsilon < \frac{1}{2}$, such that $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p-\varepsilon)}$. The proof can then be completed by setting $|\chi\rangle = |\chi(p)\rangle$ and $|\omega\rangle = |\chi(p-\varepsilon)\rangle$. We now present a construction of such a set I .

First consider the case where there are only two equalities, i.e., $\Delta_{\psi,\varphi} = \{k_1, k_2\}$, where $1 < k_1 < k_2 < n-1$ and $k_2 - k_1 \geq 2$. Since $\alpha_1 \neq \beta_1$, it cannot be the case that *both* $\alpha_1 = \alpha_{k_1}$ and $\beta_1 = \beta_{k_1}$: if it is true then the k_1^{th} inequality in the majorization is also strict and $k_1 \notin \Delta_{\psi,\varphi}$, which contradicts our assumption. Hence, $\alpha_1 > \alpha_{k_1}$ or $\beta_1 > \beta_{k_1}$ or both. Similarly, one can argue that **(i)** since $k_1 \in \Delta_{\psi,\varphi}$, and $k_1 + 1 \notin \Delta_{\psi,\varphi}$, *both* $\alpha_{k_1+1} = \alpha_{k_2}$ and $\beta_{k_1+1} = \beta_{k_2}$ cannot be true, and **(ii)** since $k_2 \in \Delta_{\psi,\varphi}$, and $k_1 + 1 \notin \Delta_{\psi,\varphi}$, *both* $\alpha_{k_2+1} = \alpha_n$ and $\beta_{k_2+1} = \beta_n$ cannot be true. Now set $I = (\frac{1}{2}, a)$, where

$$a = \min \left\{ q_1 * \frac{\alpha_1}{\alpha_1 + \alpha_{k_1}}, q_2 * \frac{\beta_1}{\beta_1 + \beta_{k_1}}, q_3 * \frac{\alpha_{k_1+1}}{\alpha_{k_1+1} + \alpha_{k_2}}, q_4 * \frac{\beta_{k_1+1}}{\beta_{k_1+1} + \beta_{k_2}}, \right. \\ \left. q_5 * \frac{\alpha_{k_2+1}}{\alpha_{k_2+1} + \alpha_n}, q_6 * \frac{\beta_{k_2+1}}{\beta_{k_2+1} + \beta_n} \right\}, \quad (4)$$

and $q_i = 2$ if its accompanying multiplicative term equals $\frac{1}{2}$, otherwise $q_i = 1$. Thus, if $\alpha_1 = \alpha_{k_1}$, then $q_1 = 2$ and the first term, $q_1 * \frac{\alpha_1}{\alpha_1 + \alpha_{k_1}} = 1$, plays no role in determining the value of a ; otherwise, if $\alpha_1 > \alpha_{k_1}$, then $q_1 = 1$ and the first term, $\frac{1}{2} < q_1 * \frac{\alpha_1}{\alpha_1 + \alpha_{k_1}} < 1$, can potentially determine a . By construction, $\frac{1}{2} < a < 1$, and hence, I is nonempty. The motivation of defining a as above is that by restricting $p \in (\frac{1}{2}, a)$, it enforces a *partial ordering* of the Schmidt coefficients of $|\varphi\rangle \otimes |\chi(p)\rangle$ and $|\psi\rangle \otimes |\chi(p)\rangle$. For example, if $\beta_1 > \beta_{k_1}$ then from (4) it follows that $p\beta_{k_1} < (1-p)\beta_1$, and hence, in the ordering of the Schmidt coefficients of $|\varphi\rangle \otimes |\chi(p)\rangle$, $(1-p)\beta_1$ will appear before $p\beta_{k_1}$.

Next, we sketch how to show $\lambda_{\psi \otimes \chi(p)} \sqsubseteq \lambda_{\varphi \otimes \chi(p)}$ for all $p \in F \cap I$, except at most $2n - 1$ discrete values. If in the majorization relationship of $\lambda_{\psi \otimes \chi(p)} \prec \lambda_{\varphi \otimes \chi(p)}$ one of the inequalities (among the

$2n - 1$ nontrivial inequalities) is an equality, then following arguments similar to those used in the proof of Theorem 1 and using the partial ordering of Schmidt coefficients enforced by the selection of I (see (4)), one can show that either **(i)** the equality determines a value of p (hence, there are at most $2n - 1$ discrete values of p where any such equality can exist), or **(ii)** the equality is a *benign* identity with one of the following forms

$$p \sum_{i=1}^{k_j} \alpha_i + (1-p) \sum_{i=1}^{k_j} \alpha_i = p \sum_{i=1}^{k_j} \beta_i + (1-p) \sum_{i=1}^{k_j} \beta_i, \quad (5)$$

where $j \in \{1, 2\}$. The reason identities like (5) are benign for our purposes is that when p is substituted by $p - \varepsilon$ on the right hand side, then the identity still remains an equality:

$$p \sum_{i=1}^{k_1} \alpha_i + (1-p) \sum_{i=1}^{k_1} \alpha_i = (p - \varepsilon) \sum_{i=1}^{k_1} \beta_i + (1 - p + \varepsilon) \sum_{i=1}^{k_1} \beta_i.$$

To prove the above claim, consider an equality in the majorization relationship, which as discussed in the proof of Theorem 1 (see (3)), can be written as:

$$\left(\sum_{j=1}^x \alpha_j - \sum_{j=1}^y \alpha_j - \sum_{j=1}^s \beta_j + \sum_{j=1}^t \beta_j \right) p = \sum_{j=1}^t \beta_j - \sum_{j=1}^y \alpha_j, \quad (6)$$

where $x + y = s + t$, $x \geq y$, and $s \geq t$. Equation (6) is an equivalence if and only if the following two conditions are simultaneously satisfied: (i) $\sum_{j=1}^t \beta_j = \sum_{j=1}^y \alpha_j$, which is true only if $t = y \in \{0, k_1, k_2\}$, or

if $y > t$; and (ii) $\sum_{j=1}^x \alpha_j = \sum_{j=1}^s \beta_j$, which is true only if $x = s \in \{k_1, k_2, n\}$, or if $x > s$. The benign identity cases occur if $x = y = s = t = k_j$, $j \in \{1, 2\}$. Let us then consider *all the other potentially feasible cases* and show that they are all *impossible*: **(i)** If $(y > t \text{ and } x \geq s)$ or $(y \geq t \text{ and } x > s)$ then we reach the contradiction that $x + y > s + t$; **(ii)** If $(y = t = 0)$ and $(x = s \in \{k_1, k_2, n\})$: this implies that $(1-p)\alpha_1 < p\alpha_{k_1}$ and $(1-p)\beta_1 < p\beta_{k_1}$, which contradicts the fact that $p \in (\frac{1}{2}, a)$ (see (4)); **(ii)** If $(y = t = k_1)$ and $(x = s \in \{k_2, n\})$: this implies that $(1-p)\alpha_{k_1+1} < p\alpha_{k_2}$ and $(1-p)\beta_{k_1+1} < p\beta_{k_2}$, which again contradicts the construction introduced in (4); **(iii)** If $(y = t = k_2)$ and $(x = s = n)$: this implies that $(1-p)\alpha_{k_2+1} < p\alpha_n$ and $(1-p)\beta_{k_2+1} < p\beta_n$, which again contradicts the construction introduced in (4).

In general, when $\Delta_{\psi, \varphi} = \{k_1, k_2, \dots, k_\ell\}$, where $1 < k_1 < \dots < k_\ell < n - 1$ and $k_{i+1} - k_i \geq 2$,

then one can show the above results for $I = (\frac{1}{2}, a)$, where

$$a = \min \left\{ q_1 * \frac{\alpha_1}{\alpha_1 + \alpha_{k_1}}, q_2 * \frac{\beta_1}{\beta_1 + \beta_{k_1}}, \dots, q_{2\ell-1} * \frac{\alpha_{k_{\ell-1}+1}}{\alpha_{k_{\ell-1}+1} + \alpha_{k_\ell}}, \right. \\ \left. q_{2\ell} * \frac{\beta_{k_{\ell-1}+1}}{\beta_{k_{\ell-1}+1} + \beta_{k_\ell}}, q_{2\ell+1} * \frac{\alpha_{k_\ell+1}}{\alpha_{k_\ell+1} + \alpha_n}, q_{2(\ell+1)} * \frac{\beta_{k_\ell+1}}{\beta_{k_\ell+1} + \beta_n} \right\},$$

and q_i 's are chosen as in (4). ■

We next show that for *certain equality patterns* in the majorization relation, *partial recovery* with the help of 2×2 states (or even 3×3 states) is *not always possible*.

Lemma 1 *If $\alpha_1 = \beta_1$ or $\alpha_n = \beta_n$, then recovery is not possible with 2×2 auxiliary states. Also, if both relations $\alpha_1 = \beta_1$ and $\alpha_n = \beta_n$ hold then there is no recovery even with 3×3 auxiliary states*

Proof: First assume that $\alpha_1 = \beta_1$ or $\alpha_n = \beta_n$. Suppose, by contradiction, there are 2×2 states $|\chi\rangle$ and $|\omega\rangle$ such that $|\psi\rangle \otimes |\chi\rangle \longrightarrow |\varphi\rangle \otimes |\omega\rangle$ and $E(|\omega\rangle) > E(|\chi\rangle)$. Let $\lambda_\chi = (p, 1-p)$ and $\lambda_\omega = (q, 1-q)$ be the vector of eigenvectors of $|\chi\rangle$ and $|\omega\rangle$ with $p, q > \frac{1}{2}$. The condition $E(|\omega\rangle) > E(|\chi\rangle)$ implies that $q < p$. The relation $\lambda_{\psi \otimes \chi} \prec \lambda_{\varphi \otimes \omega}$ implies that $\alpha_1 p \leq \beta_1 q$ and $1 - \alpha_n(1-p) \leq 1 - \beta_n(1-q)$. So if $\alpha_1 = \beta_1$ or $\alpha_n = \beta_n$ then $p \leq q$ and $E(|\omega\rangle) \leq E(|\chi\rangle)$, which is a contradiction.

Now suppose that $\alpha_1 = \beta_1$ and $\alpha_n = \beta_n$. Let us assume, by contradiction, that there are 3×3 recovery states $|\chi\rangle$ and $|\omega\rangle$ with eigenvalue vectors $\lambda_\chi = (p, q, 1-p-q)$ and $\lambda_\omega = (p', q', 1-p'-q')$ with $p \geq q \geq 1-p-q$ and $p' \geq q' \geq 1-p'-q'$. Then $\lambda_{\psi \otimes \chi} \prec \lambda_{\varphi \otimes \omega}$ implies that $\alpha_1 p \leq \beta_1 p'$ and $1 - \alpha_n(1-p-q) \leq 1 - \beta_n(1-p'-q')$. Thus $p \leq p'$ and $p+q \leq p'+q'$. Therefore, $\lambda_\chi \prec \lambda_\omega$ and $|\chi\rangle \longrightarrow |\omega\rangle$. The last relation implies that $E(|\omega\rangle) \leq E(|\chi\rangle)$ (see [5]), which is a contradiction. ■

The preceding result shows that if $\alpha_1 = \beta_1$ then any auxiliary state for partial recovery should be at least a 3×3 state. Here we show that such minimum-dimension auxiliary states can exist.

Theorem 3 *If $\Delta_{\psi, \varphi} = \{1\}$ then there are 3×3 states $|\chi\rangle$ and $|\omega\rangle$ such that $|\psi\rangle \otimes |\chi\rangle \longrightarrow |\varphi\rangle \otimes |\omega\rangle$ and $E(|\omega\rangle) > E(|\chi\rangle)$.*

Proof: Let $|\chi(p, q)\rangle$ be a 3×3 state with $\lambda_{\chi(p, q)} = (p, q, 1-p-q)$, where $p \geq q \geq 1-p-q \geq 0$. The goal is to find the state $|\omega\rangle$ of the form $|\chi(p, q - \varepsilon)\rangle$, for some $\varepsilon > 0$. Our approach will be similar to that introduced in the proof of Theorem 2. In particular, it is sufficient to show that there exists a region in the plane, $R = \{(p, q) | p \geq q \geq 1-p-q \geq 0\}$, with nonzero area such that $\lambda_{\psi \otimes \chi(p, q)} \sqsubseteq \lambda_{\varphi \otimes \chi(p, q)}$ for almost all $(p, q) \in R$, and the set of points where it is violated has measure zero. Thus for almost all the points in R , each inequality in the majorization relationship $\lambda_{\psi \otimes \chi(p, q)} \prec \lambda_{\varphi \otimes \chi(p, q)}$ is either (i) a “benign” identity;

that is, the equality holds even if on the right hand side, (p, q) is perturbed to $(p, q - \varepsilon)$, for any $0 < \varepsilon$, or **(ii)** is a *strict* inequality. The measure-theoretic arguments used in the proofs of Theorems 1 and 2 will guarantee that there is an $\varepsilon > 0$, such that $\lambda_{\psi \otimes \chi(p, q)} \prec \lambda_{\varphi \otimes \chi(p, q - \varepsilon)}$. The proof can then be completed by setting $|\chi\rangle = |\chi(p)\rangle$ and $|\omega\rangle = |\chi(p - \varepsilon)\rangle$.

We now present a construction of such a set R . First note that, since $2 \notin \Delta_{\psi, \varphi}$, hence $\alpha_1 > \alpha_2$. Also note that if $\alpha_2 = \alpha_n$ then $\beta_2 > \beta_n$. Therefore, we have to consider two case: **(i)** $\alpha_1 > \alpha_2 > \alpha_n$; and **(ii)** $\alpha_1 > \alpha_2 = \alpha_n$ and $\beta_2 > \beta_n$. It is easy to check that in the case **(ii)** we have $\alpha_2 > \beta_n$.

To define R , we choose the parameters p and q such that $p \geq q \geq 1 - p - q \geq 0$ and they satisfy the following conditions. case **(i)**:

$$q \alpha_1 < p \alpha_2, \quad p \alpha_n < (1 - p - q) \alpha_2; \quad (7)$$

or, case **(ii)**:

$$q \alpha_1 < p \alpha_2, \quad p \beta_n < (1 - p - q) \beta_1. \quad (7')$$

Note that the systems (7) and (7') imply $q \alpha_n < (1 - p - q) \alpha_2$ and $q \beta_n < (1 - p - q) \beta_2$, respectively. So the region R is either given by $p \geq q \geq 1 - p - q \geq 0$, and

$$q < \frac{\alpha_2}{\alpha_1} p, \quad q < 1 - \frac{\alpha_2 + \alpha_n}{\alpha_2} p, \quad (8)$$

or by

$$q < \frac{\alpha_2}{\alpha_1} p, \quad q < 1 - \frac{\beta_1 + \beta_n}{\beta_1} p. \quad (8')$$

In both cases (i.e., (8) or (8')), R defines a non-empty triangular region in the (p, q) -plane (see Figure 1).

Now for any $(p, q) \in R$, one can show that if any of the $3n$ inequalities in the majorization relationship $\lambda_{\psi \otimes \chi(p, q)} \prec \lambda_{\varphi \otimes \chi(p, q)}$ is an equality, then it belongs to one of the two following cases. Case**(i)**: It is a *non identical* equality, i.e., the set of (p, q) that satisfies it defines a line in (p, q) plane, and hence comprises a measure zero set. Since, the total number of such non identical equalities is at most $3n$, the set of all points in R where there might be a non-identical equality is of measure zero. Case**(ii)**: It is a *benign* identity of the form :

$$p + q \alpha_1 + (1 - p - q) \alpha_1 = p + q \beta_1 + (1 - p - q) \beta_1.$$

Therefore, R satisfies the property that $\lambda_{\psi \otimes \chi(p, q)} \sqsubseteq \lambda_{\varphi \otimes \chi(p, q)}$ for almost all $(p, q) \in R$. ■

Let $\eta_{\psi, \varphi}$ be the size of the longest block of consecutive equalities in the majorization relationship of $|\psi\rangle$ and $|\varphi\rangle$; i.e., $\eta_{\psi, \varphi}$ is the largest integer m such that $\{j, j + 1, \dots, j + m - 1\} \subseteq \Delta_{\psi, \varphi}$. Our final

theorem is a general construction which shows that if $\alpha_n \neq \beta_n$, then partial recovery of entanglement is always possible using auxiliary states of dimension $k = \eta_{\psi,\varphi} + 2$. Thus, if $\alpha_n \neq \beta_n$, then *genuine* partial recovery is *always possible*: since $\alpha_n \neq \beta_n$, $\eta_{\psi,\varphi} \leq (n - 3)$ and **hence** $k < n$.

Theorem 4 *If $n - 1 \notin \Delta_{\psi,\varphi}$, then there are $k \times k$ states $|\chi\rangle$ and $|\omega\rangle$ such that $k = \eta_{\psi,\varphi} + 2$, $|\psi\rangle \otimes |\chi\rangle \longrightarrow |\varphi\rangle \otimes |\omega\rangle$, and $E(|\omega\rangle) > E(|\chi\rangle)$.*

Proof: We provide the proof for a special case. The proof can be easily extended to cover any general case .

Suppose that $\Delta_{\psi,\varphi} = \{2, 3, 5\}$. So $\eta_{\psi,\varphi} = 2$. Let $|\chi(p_1, p_2, p_3)\rangle$ be a 4×4 state with $\lambda_{\chi(p_1, p_2, p_3)} = (p_1, p_2, p_3, p_4)$, where $p_4 = 1 - p_1 - p_2 - p_3$ and $p_1 \geq p_2 \geq p_3 \geq p_4 \geq 0$. Let

$$S = \{(p_1, p_2, p_3) \in \mathbb{R}^3 : p_1 \geq p_2 \geq p_3 \geq 1 - p_1 - p_2 - p_3 \geq 0\}.$$

Then S is the subset of $\mathbb{R}^3$ bounded by the following polyhedral:

$$\left. \begin{aligned} p_1 &\geq 0, & p_2 &\geq 0, & p_3 &\geq 0, \\ p_1 &\geq p_2, & p_2 &\geq p_3, \\ p_1 + p_2 + p_3 &\leq 1, \\ p_1 + p_2 + 2p_3 &\geq 1. \end{aligned} \right\} \quad (9)$$

The goal is to find the state $|\omega\rangle$ of the form $|\chi(p_1, p_2, p_3 - \varepsilon)\rangle$, for some $\varepsilon > 0$. As in the proof of Theorem 3, it suffices to show that there is a region $R \subseteq S$ such that $\lambda_{\psi \otimes \chi(p_1, p_2, p_3)} \sqsubseteq \lambda_{\varphi \otimes \chi(p_1, p_2, p_3)}$ for almost all $(p_1, p_2, p_3) \in R$, and the set of points where it is violated has measure zero. The following conditions guarantee the only identical equalities among majorization relations of $\lambda_{\psi \otimes \chi(p_1, p_2, p_3)} \prec \lambda_{\varphi \otimes \chi(p_1, p_2, p_3)}$ are benign.

$$\left. \begin{aligned} \frac{p_3}{p_1} &< \min \left(\frac{\alpha_4}{\alpha_2}, \frac{\beta_4}{\beta_2} \right) \\ \frac{p_4}{p_1} &> \min \left(\frac{\alpha_5}{\alpha_2}, \frac{\beta_5}{\beta_2} \right) \\ \frac{p_3}{p_2} &< \min \left(\frac{\alpha_4}{\alpha_3}, \frac{\beta_4}{\beta_3} \right) \\ \frac{p_4}{p_2} &> \min \left(\frac{\alpha_5}{\alpha_3}, \frac{\beta_5}{\beta_3} \right) \\ \frac{p_3}{p_1} &< \min \left(\frac{\alpha_5}{\alpha_6}, \frac{\beta_5}{\beta_6} \right) \\ \frac{p_4}{p_1} &> \min \left(\frac{\alpha_n}{\alpha_6}, \frac{\beta_n}{\beta_6} \right) \\ \frac{p_4}{p_3} &> \min \left(\frac{\alpha_n}{\alpha_6}, \frac{\beta_n}{\beta_6} \right) \end{aligned} \right\} \quad (10)$$

The assumption $\Delta_{\psi,\varphi} = \{2, 3, 5\}$ implies that among each pair of numbers in the right-hand side of the above inequalities at least one is less than 1. Also note that for any point $(p_1, p_2, p_3) \in S$, we have

$$\frac{p_3}{p_1} \leq \frac{p_3}{p_2} \quad \text{and} \quad \frac{p_4}{p_1} \leq \frac{p_4}{p_3}.$$

So there are real numbers s and t such that $0 < s, t < 1$ and if

$$\frac{p_3}{p_2} < t, \tag{11}$$

$$\frac{p_4}{p_1} > s, \tag{12}$$

then all conditions (10) are satisfied. Now we show that the conditions (11) and (12) define a subset of S with nonzero measure. The condition (12) defines a half-space $(1+s)p_1 + p_2 + p_3 < 1$ which is a subset of the half-space $p_1 + p_2 + p_3 \leq 1$. So in the first step, instead of S , we consider the region

$$\left. \begin{aligned} p_1 &\geq 0, & p_2 &\geq 0, & p_3 &\geq 0, \\ p_1 &\geq p_2, & p_2 &\geq p_3, \\ (1+s)p_1 + p_2 + p_3 &< 1, \\ p_1 + p_2 + 2p_3 &\geq 1. \end{aligned} \right\} \tag{13}$$

The condition (11) defines a half-space $p_2 > \frac{1}{t} p_3$ which is a subset of the half-space $p_2 \geq p_3$ and cuts a non-empty subset R of the region defined by (9).

Like the proof of Theorem 2, for some feasible set F of a given ordering of the Schmidt coefficients of $|\phi\rangle \otimes |\chi(p_1, p_2, p_3)\rangle$ the intersection $F \cap R$ has nonzero measure. The points $(p_1, p_2, p_3) \in F \cap R$ for which there are nonidentical identities among majorization relations of $\lambda_{\psi \otimes \chi(p_1, p_2, p_3)} \prec \lambda_{\varphi \otimes \chi(p_1, p_2, p_3)}$ belong to a *finite* set of hyper-planes; i.e., a measure zero set. Also the conditions (10) implies that all the other identities among coefficients are benign. So there is an $\varepsilon > 0$ such that $\lambda_{\psi \otimes \chi(p_1, p_2, p_3)} \prec \lambda_{\varphi \otimes \chi(p_1, p_2, p_3 - \varepsilon)}$.

In summary, we have shown the following results:

1. If all the inequalities of the majorization conditions (1) are strict then recovery with an auxiliary entangled state in 2×2 is always possible. Note that *for a randomly picked pair of comparable states the majorization inequalities are strict with probability one*.
2. If $\alpha_1 \neq \beta_1$ and $\alpha_n \neq \beta_n$ and in the majorization conditions there are no two consecutive equalities, then recovery with an auxiliary entangled state in 2×2 is possible.
3. If $\alpha_1 = \beta_1$ or $\alpha_n = \beta_n$ then recovery with a 2×2 auxiliary state is not possible; and if $\alpha_1 = \beta_1$ and $\alpha_n = \beta_n$ then recovery with a 3×3 auxiliary state is not possible.

4. If $\alpha_n \neq \beta_n$ and the number of consecutive equalities in the majorization conditions (1) are m , then recovery is always possible using $(m+2) \times (m+2)$ auxiliary states. Hence if $\alpha_n \neq \beta_n$, in the worst case, a recovery by $(n-1) \times (n-1)$ states is achievable; i.e., a genuine recovery is always possible.

Thus, we have shown a nontrivial recovery is always possible except for the special case where $\alpha_n = \beta_n$; whether recovery is still possible for this special case is left as an open problem. There are many other open questions that might be of interest. For example, for a given pair of comparable states, one may ask what is the maximum entanglement that can be recovered. Similarly, can one recover more entanglement by increasing the dimension of the auxiliary entangled states? For example, we show that for almost all comparable states, 2×2 auxiliary states are sufficient to implement partial recovery; however, can one have more recovery of entanglement if the dimension of the auxiliary state is increased? We hope that the results of the present work will lead to a better understanding of the subtleties involved in local entanglement manipulation in higher dimensions.

References

- [1] E. Schrodinger, *Naturwissenschaften* **23**, 807 (1935); **23**, 823 (1935); **23**, 844 (1935); For a review see: M. B. Plenio and V. Vedral, *Cont. Phys.* **39**, 431 (1998).
- [2] C. H. Bennett, *Physics Today* **48**, 24 (1995).
- [3] J. Preskill, *Proc. Roy. Soc. A: Math., Phys. and Eng.* **454**, 469 (1998);
- [4] C. H. Bennett, H.J. Bernstein, S. Popescu, and B. Schumacher, *Phys. Rev. A* **53**, 2046 (1996); C. H. Bennett, D. P. DiVincenzo, J. A. Smolin and W. K. Wootters, *Phys. Rev. A* **54**, 3814 (1997).
- [5] M. A. Nielsen, *Phys. Rev. Lett.* **83**, 436 (1999).
- [6] H. K. Lo and S. Popescu, “Concentrating entanglement by local actions beyond mean values”, *quant-ph/9707038*; G. Vidal, *Phys. Rev. Lett.* **83**, 1046 (1999); D. Jonathan and M. B. Plenio *Phys.Rev.Lett.* **83**, 3566 (1999); G. Vidal, D. Jonathan and M. A. Nielsen *Phys. Rev. A* **62**, 012304 (2000).
- [7] S. Bandyopadhyay, V. P. Roychowdhury, U. Sen, “A classification of incomparable states”, *quant-ph/0103131*
- [8] F. Morikoshi, *Phys.Rev.Lett.* **84**, 3189 (2000).




Figure 1: The shaded region is the solution set of the system (13).